

**The Deafening Silence is
Killing Us
EUA COVID-19 Vaccines**

August 14, 2022

Dr. Marian Laderoute

hervk102@substack.com

Globally despite clear evidence that the COVID-19 vaccines no longer provide benefit but are killing recipients at about 1/100 [data released by the Office for National Statistics, UK] the regulatory authorities worldwide have not rescinded their EUA approvals.

This is true of the USA, the WHO, Canada, and even the UK where the raw data were released on July 6, 2022.

FDA may issue an EUA when certain criteria are met, which includes that there are no adequate, approved, and available alternatives. In addition, the FDA decision is based on the totality of the scientific evidence available showing that the product may be effective to prevent COVID-19 during the COVID-19 pandemic and that the known and potential benefits of the product outweigh the known and potential risks of the product. All of these criteria must be met to allow for the product to be used during the COVID-19 pandemic.

USA

On July 6, 2022 the UK Office for National Statistics released its report

<https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/deathsinvolvingcovid19byvaccinationstatusengland/deathsoccurringbetween1january2021and31may2022>

which indicated for England

Mass vaccination saved 2110 lives from COVID-19 deaths but caused 423,337 vaccine toxicity deaths (201 additional deaths per life saved by mass vaccination of about 57 million people). How does this indicate **the benefits of COVID-19 vaccines continue to outweigh the risks of disease????**

https://www.who.int/news-room/feature-stories/detail/who-can-take-the-pfizer-biontech-covid-19--vaccine-what-you-need-to-know

The Pfizer BioNTech (BNT162b2) COVID-19 vaccine: What you need to know

10 June 2022

Updated 10 June 2022, to ensure consistency of formatting

The WHO Strategic Advisory Group of Experts on Immunization (SAGE) has issued interim recommendations for the use of the Pfizer BioNTech (BNT162b2) vaccine against COVID-19. This article provides a summary of those interim recommendations; you may access the full guidance document [here](#).

Here is what you need to know.

According to SAGE, the Pfizer-BioNTech COVID-19 mRNA vaccine is safe and effective. The priority is to start vaccinating health workers at high risk of exposure, followed by older adults, before immunizing the rest of the population.

Who can take the vaccine?

The vaccine is safe and effective for all individuals aged 5 and above. In line with the [WHO Prioritization Roadmap](#) and the [WHO Values Framework](#), older adults, health workers and immunocompromised persons should be prioritised.

The Pfizer vaccine can be offered to individuals who have had COVID-19 in the past, but individuals may wish to delay vaccination for 3 months following the infection

Is it safe?

The Global Advisory Committee on Vaccine Safety (GACVS), a group of experts that provides independent and authoritative guidance to WHO on the topic of safe vaccine use, receives and assesses reports of suspected safety events of potentially international impact. In October 2021, the GACVS COVID-19 subcommittee concluded that the mRNA COVID-19 vaccines have clear benefits in all age groups in reducing hospitalizations and deaths due to COVID-19.

How efficacious is the vaccine?

The Pfizer BioNTech vaccine against COVID-19 has an efficacy of 95% against symptomatic SARS-CoV-2 infection.

UPDATED **10 June 2022**
WHO



MENU ▾

[Canada.ca](#) > [Coronavirus disease \(COVID-19\)](#)

Vaccines for COVID-19

Free vaccines are available to everyone in Canada.

[Get a vaccine or booster](#)

CANADA



[COVID-19 vaccines for children and youth](#)

Your vaccination

- [Booster doses](#)
- [Proof of vaccination, QR code](#)
- [Effectiveness and benefits](#)
- [Vaccinated travellers entering Canada](#)

Vaccine safety

- [Vaccine safety and side effects](#)
- [Reported side effects](#)
- [Clinical trials](#)
- [How vaccines are authorized](#)
- [Monitoring after authorization](#)

Authorized vaccines

- [AstraZeneca Vaxzevria](#)
- [Janssen Jcovden \(Johnson & Johnson\)](#)
- [Medicago Covifenz](#)
- [Moderna Spikevax](#)
- [Novavax Nuvaxovid](#)
- [Pfizer-BioNTech Comirnaty](#)
- [Other vaccines under review](#)
- [Canada's vaccine supply and donation](#)

- Evidence indicates that the benefits of COVID-19 vaccines continue to outweigh the risks of the disease.

Canada does not publically reveal the raw counts of COVID-19 deaths, non-COVID-19 deaths and all-cause deaths by vaccination status. However, the Office for National Statistics in England does. The latest was published July 6, 2022.

<https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/deathsinvolvingcovid19byvaccinationstatusengland/deathsoccurringbetween1january2021and31may2022>

FAILED All-cause Mortality Test

From the start of the mass vaccination in England until the end of May 2022, for each life saved from a COVID-19 death, the mass vaccination campaign caused 201 additional vaccine injury deaths.

Summary: From January 2021 to May 2022 the COVID-19 vaccines saved 2110 people from COVID-19 deaths, but the vaccines were associated with 423,337 additional non-COVID-19 deaths with a net of 421,227 all cause deaths, in England.

| AGE | COVID-19 Deaths | | | NON-COVID Deaths | | | All-CAUSE | | |
|---------|-----------------|---------|-------|------------------|---------|--------|-----------|---------|--------|
| | Unvax | Evervax | E-U | Unvax | Evervax | E-U | Unvax | Evervax | E-U |
| | 90+ | 7191 | 9336 | 2145 | 11829 | 116270 | 104441 | 19020 | 125606 |
| 85-89 | 6437 | 7870 | 1433 | 9707 | 94963 | 85256 | 16144 | 102833 | 86689 |
| 80-84 | 5841 | 6620 | 779 | 8887 | 84170 | 75283 | 14728 | 90790 | 76062 |
| 75-79 | 5044 | 4712 | -332 | 8515 | 67288 | 58773 | 13559 | 72000 | 58441 |
| 70-74 | 4194 | 3031 | -1163 | 8090 | 49631 | 41541 | 12284 | 52662 | 40378 |
| 65-69 | 3051 | 1776 | -1275 | 6303 | 29422 | 23119 | 9354 | 31198 | 21844 |
| 60-64 | 2425 | 1162 | -1263 | 5135 | 20397 | 15262 | 7560 | 21559 | 13999 |
| 55-59 | 1626 | 754 | -872 | 4084 | 13903 | 9819 | 5710 | 14657 | 8947 |
| 50-54 | 1069 | 407 | -662 | 3014 | 8471 | 5457 | 4083 | 8878 | 4795 |
| 45-49 | 597 | 223 | -374 | 1965 | 4416 | 2451 | 2562 | 4639 | 2077 |
| 40-44 | 299 | 121 | -178 | 1225 | 2480 | 1255 | 1524 | 2601 | 1077 |
| 35-39 | 238 | 71 | -167 | 903 | 1410 | 507 | 1141 | 1481 | 340 |
| 30-34 | 129 | 40 | -89 | 649 | 886 | 237 | 778 | 926 | 148 |
| 25-29 | 68 | 26 | -42 | 525 | 577 | 52 | 593 | 603 | 10 |
| 20-24 | 43 | 16 | -27 | 335 | 397 | 62 | 378 | 413 | 35 |
| 15-19 | 24 | 7 | -17 | 265 | 225 | -40 | 289 | 232 | -57 |
| 10-14 | 9 | 3 | -6 | 175 | 37 | -138 | 184 | 40 | -144 |
| Total | 74460 | 38285 | 36175 | 566549 | 71606 | 494943 | 641009 | 109891 | 531118 |
| Summary | | | -2110 | | | 423337 | | | 421227 |

| At least one dose analysis | C19 DEATHS | Non-C19 Deaths | All-Cause Deaths |
|---|------------|----------------|------------------|
| Actual % of total deaths in the unvax | 51.42% | 12.64% | 17.14% |
| Expected % of total deaths in the unvax | 23.95% | 23.95% | 23.95% |
| Favors | Vaxed | Unvaxed | Unvaxed |
| | 0.58 | 0.42 | 0.57 |

For each life saved from a SARS-COV-2 infection, vaccination with 1-3 doses caused 201 vaccine injury deaths.

Mass vaccination saved 2110 lives from COVID-19 deaths, but caused 423,337 vaccine toxicity deaths (201 additional deaths per life saved by mass vaccination of about 57 million people) indicates the benefits of COVID-19 vaccines continue to outweigh the risks of disease????

<https://www.canada.ca/en/public-health/services/diseases/coronavirus-disease-covid-19/vaccines/safety-side-effects.html>

From
December 2020
to May 29,
2022

423337 excess non-Covid-19
deaths in 44,386,966 ever
vaxed= **1/105 ever vaxed.**

423337 excess non-Covid-19
Deaths in 32,770,364
3rd dose vaxed = **1/77 3rd dose vaxed.**

Table 5: Provisional cumulative COVID-19 vaccine uptake by age in England

| NATIONAL | People in NIMS cohort | Vaccinated with at least 1 dose | | Vaccinated with at least 2 doses | | Vaccinated with at least 3 doses | |
|----------------|-----------------------|---------------------------------|------------------|----------------------------------|------------------|----------------------------------|------------------|
| | | Number vaccinated | % vaccine uptake | Number vaccinated | % vaccine uptake | Number vaccinated | % vaccine uptake |
| Over 80 | 2,934,025 | 2,807,002 | 95.7 | 2,789,675 | 95.1 | 2,712,787 | 92.5 |
| 75 to under 80 | 2,337,404 | 2,239,354 | 95.8 | 2,224,226 | 95.2 | 2,169,105 | 92.8 |
| 70 to under 75 | 2,774,515 | 2,618,019 | 94.4 | 2,595,524 | 93.5 | 2,506,954 | 90.4 |
| 65 to under 70 | 2,983,896 | 2,756,680 | 92.4 | 2,725,075 | 91.3 | 2,581,751 | 86.5 |
| 60 to under 65 | 3,615,299 | 3,282,399 | 90.8 | 3,236,637 | 89.5 | 2,976,508 | 82.3 |
| 55 to under 60 | 4,170,017 | 3,715,500 | 89.1 | 3,653,334 | 87.6 | 3,273,048 | 78.5 |
| 50 to under 55 | 4,246,640 | 3,679,320 | 86.6 | 3,602,866 | 84.8 | 3,122,431 | 73.5 |
| 45 to under 50 | 3,957,299 | 3,248,281 | 82.1 | 3,157,889 | 79.8 | 2,580,202 | 65.2 |
| 40 to under 45 | 4,331,967 | 3,346,067 | 77.2 | 3,222,198 | 74.4 | 2,473,682 | 57.1 |
| 35 to under 40 | 4,670,380 | 3,397,463 | 72.7 | 3,234,521 | 69.3 | 2,301,945 | 49.3 |
| 30 to under 35 | 4,887,611 | 3,410,849 | 69.8 | 3,202,158 | 65.5 | 2,119,675 | 43.4 |
| 25 to under 30 | 4,557,546 | 3,117,575 | 68.4 | 2,886,123 | 63.3 | 1,788,762 | 39.2 |
| 20 to under 25 | 3,925,321 | 2,828,843 | 72.1 | 2,556,321 | 65.1 | 1,502,563 | 38.3 |
| 18 to under 20 | 1,387,814 | 1,018,119 | 73.4 | 893,590 | 64.4 | 467,420 | 33.7 |
| 16 to under 18 | 1,388,641 | 911,375 | 65.6 | 686,566 | 49.4 | 177,843 | 12.8 |
| 12 to under 16 | 2,937,494 | 1,577,277 | 53.7 | 1,011,656 | 34.4 | 15,458 | 0.5 |
| 5 to under 12 | 5,080,285 | 431,969 | 8.5 | 16,988 | 0.3 | 23 | 0.0 |
| Under 5 | 3,067,754 | 114 | 0.0 | 11 | 0.0 | 2 | 0.0 |
| Total* | 63,283,908 | 44,386,966 | 70.1 | 41,695,721 | 65.9 | 32,770,364 | 51.8 |

England Population Statistics Vaccinated by AGE versus Total https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1080640/Weekly_Flu_and_COVID-19_report_w22.pdf

<https://www.ons.gov.uk/populationandmigration/population>

Population estimates - Office for National Statistics

On Census Day, 21 March 2021, the size of the usual resident population in England and Wales was 59,597,300 (56,489,800 in England and 3,107,500 in Wales); ...

The numbers don't
match.

Decision

Regulatory approval of Pfizer/BioNTech vaccine for COVID-19

Information for healthcare professionals and the public about the Pfizer/BioNTech vaccine.

From: [Medicines and Healthcare products Regulatory Agency](#)

Published 2 December 2020

Last updated 7 July 2022 — [See all updates](#)

No mention of non-COVID-19 deaths due to COVID-19 vaccine toxicity 1/105 for ever vaxed or 1/77 for 3rd dose vaxed. DATA available July 6, 2022 Office for National Statistics (UK).

4. Possible side effects

Like all vaccines, Comirnaty can cause side effects, although not everybody gets them.

Very common side effects: may affect more than 1 in 10 people

- injection site: pain, swelling
- tiredness
- headache
- muscle pain
- chills
- joint pain
- diarrhoea
- fever

Some of these side effects were slightly more frequent in adolescents 12 to 15 years than in adults.

Common side effects: may affect up to 1 in 10 people

- injection site redness
- nausea
- vomiting

Uncommon side effects: may affect up to 1 in 100 people

- enlarged lymph nodes (more frequently observed after the booster dose)
- feeling unwell
- arm pain
- insomnia
- injection site itching
- allergic reactions such as rash or itching
- feeling weak or lack of energy/sleepy
- decreased appetite
- excessive sweating
- night sweats

Very rare side effects: may affect up to 1 in 10,000 people

- inflammation of the heart muscle (myocarditis) or inflammation of the lining outside the heart (pericarditis) which can result in breathlessness, palpitations or chest pain

Not known (cannot be estimated from the available data)

- severe allergic reaction
- extensive swelling of the vaccinated limb
- swelling of the face (swelling of the face may occur in patients who have had facial dermatological fillers)
- a skin reaction that causes red spots or patches on the skin, that may look like a target or “bulls-eye” with a dark red centre surrounded by paler red rings (erythema multiforme)
- unusual feeling in the skin, such as tingling or a crawling feeling (paraesthesia)
- decreased feeling or sensitivity, especially in the skin (hypoesthesia)

**The silence is deafening on the
lack of truth by all regulatory
authorities around the world that
the vaccines are causing deaths at
an ever-growing rate and no longer
provide much in terms of benefit.**

**There is no mention of non-COVID-19 deaths
due to COVID-19 vaccine toxicity
(at 1/105 for ever vaxed or 1/77 for 3rd dose vaxed).**

**Yet, the DATA was available worldwide on July 6, 2022,
Office for National Statistics (UK).**